

7969-199

7/8/2014

J adler  
1/47



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Ms. Christine M. Keating  
Product Registration Manager  
26 Davis Drive  
Research Triangle Park, NC 27709-3528

JUL 08 2014

Subject: Pristine Fungicide  
EPA Reg. No. 7969-199  
EPA Decision Number: 492621  
Your master and globe artichoke supplemental label submitted on June 17, 2014 to update tank mix instructions; add citrus black spot to citrus diseases; remove California restrictions; and roll in the radicchio supplemental label

Dear Ms. Keating:

The master label and supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended are acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

One copy of the labels stamped "Accepted" are enclosed for your records. If you have any questions, please contact Heather Garvie by phone at: 703-308-0034 or via email at: [garvie.heather@epa.gov](mailto:garvie.heather@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Shaja B. Joyner".

Shaja B. Joyner  
Product Manager 20  
Fungicide Branch  
Registration Division

Enclosure: Stamped labels "Accepted"

2/47



# Pristine<sup>®</sup>

**FUNGICIDE**

**ACCEPTED**  
**JUL 08 2014**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No. **7969-199**

For use in disease control and plant health in the following crops: alfalfa; Belgium endive; berries; bulb vegetables; carrot; celery; citrus fruit; cotton; cucurbit vegetables; dry beans; fruiting vegetables; globe artichoke; grape; hops; leafy greens; leafy petioles; low growing berry; oilseed crops; peanut; persimmon; pome fruit; small fruit, vine climbing; soybean; spinach; stone fruit; strawberry; tree nut; tropical fruit; and turnip greens

**Active Ingredients:**

pyraclostrobin*: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy)methyl]phenyl]methoxy-, methyl ester) . . . . .	12.8%
boscalid**: 3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)- . . . . .	25.2%
<b>Other Ingredients:</b> . . . . .	62.0%
<b>Total:</b> . . . . .	100.0%

\* 0.128 oz (0.008 lb) of pyraclostrobin in 1 oz of product  
\*\* 0.252 oz (0.0158 lb) of boscalid in 1 oz of product

EPA Reg No. 7969-199

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty,** and state-specific crop and/or use site restrictions.

**In case of an emergency endangering life or property involving this product, call day or night, 1-800-832-HELP (4357).**

**Net Contents:**

<b>FIRST AID</b>	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing.

#### Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to **Category A** on an EPA chemical-resistance category selection chart.

Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material (such as nitrile, butyl, neoprene, and/or barrier laminate)
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### Environmental Hazards

**DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

#### Groundwater Advisory

Boscalid and pyraclostrobin are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

#### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of boscalid and pyraclostrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when

rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours** for all crop uses except when performing cane tying, cane turning or cane girdling on grapes. The REI is **5 days** for treated grapes when conducting cane tying, cane turning or cane girdling.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, made of any waterproof material (such as nitrile, butyl, neoprene, and/or barrier laminate)
- Shoes plus socks

### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

(continued)

### STORAGE AND DISPOSAL (continued)

#### Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your state pesticide agency or environmental control agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

#### Container Handling

**Nonrefillable Container. DO NOT reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake (capacity ≤ 50 pounds) as follows.** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Triple rinse containers too large to shake (capacity > 50 pounds) as follows.** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### In Case of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

**Steps to be taken in case material is released or spilled:**

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

**Product Information**

This package contains **Pristine® fungicide**, a water dispersible granule (WG). The active ingredients in **Pristine** belong to two classes of fungicides, the strobilurins and anilides. Preventive applications optimize disease control resulting in improved plant health. The increase in plant health comes from the combined effect of disease control (including fungal diseases listed in Crop-specific directions), improved growth efficiency and improved stress tolerance. Overall increased plant health may result in an improvement in crop growth and crop quality as well as increased crop yields. **Pristine** is effective against pathogens resistant to other fungicides. **Pristine** has a protective effect because it inhibits spore germination. It also has a curative effect because it inhibits mycelial growth and sporulation of the fungus on the leaf surface. However, optimum disease control is achieved when **Pristine** is applied in a regularly scheduled protective spray program and is used in a rotation program with other fungicides. Because of its high specific activity and rainfastness, **Pristine** has good residual activity against target fungi.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

**Pristine** is not for use in greenhouse or transplant production.

**Sensitive Crop Precaution**

**Grape - DO NOT** use on Concord or Noiret (NY73.0136.17) due to foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related grape varieties. Use special care when applying **Pristine** to prevent contact with these sensitive varieties. Not all varieties have been thoroughly tested. Consult a BASF representative for more information concerning these sensitive grape varieties. Thoroughly rinse spray equipment, including the inside of the tank, hoses and nozzles after and before using the same equipment on grape varieties sensitive to **Pristine**.

**Blueberry (highbush and lowbush) - DO NOT** apply **Pristine** to blueberries as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other

additives or anything other than water. Mix **Pristine** with water only for applications to blueberries.

**Modes of Action**

Pyraclostrobin and boscalid, the active ingredients of **Pristine**, belong to the groups of respiration inhibitors classified by the U.S. EPA and Canada PMRA as target site of action **Group 7** and **Group 11** fungicides, respectively.

**Resistance Management**

**Pristine** contains pyraclostrobin and boscalid, a premix of a **Group 7** and a **Group 11** fungicide, and is effective against pathogens resistant to fungicides with modes of action different from those of target site **Group 7** and **Group 11**, such as dicarboximides, sterol inhibitors, benzimidazoles, or phenylamides. **Pristine** is also effective against certain pathogens with resistance to **Group 11** fungicides, such as pyraclostrobin, azoxystrobin, trifloxystrobin, or kresoxim-methyl. However, fungal isolates resistant to **Group 7** or **Group 11** fungicides may eventually dominate the fungal population if **Group 7** or **Group 11** fungicides are used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species, especially if resistance to either **Group 7** or **Group 11** fungicides is already present in the pathogen population. This may result in reduction of disease control by **Pristine** or other **Group 7** or **Group 11** fungicides. To maintain the performance of **Pristine** in the field, **DO NOT** exceed the specified number of applications of **Pristine** and the total number of applications of **Pristine** per season stated in **Restrictions and Limitations** and **Crop-specific Use Requirements**. Adhere to the label instructions regarding the sequential use of **Pristine** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

**Resistance Management Advisory**

The following instructions can delay the development of fungicide resistance:

1. **Tank mixtures - Pristine** provides more effective resistance management of most of its target pathogens, because it is a premix of two fungicides with different modes of action. If **Pristine** is used in tank mixtures with fungicides from different target site of action groups that are registered/permitted for the same use and that are effective against the pathogens of concern, use at least the minimum labeled rates of each fungicide in the tank mix.
2. **IPM - Integrate Pristine** into an overall disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or BASF representative for additional IPM strategies established for your area. **Pristine** may be used in agricultural extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.

3. **Monitoring** - Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.

4. **Reporting** - If a **Group 7** or **Group 11** target site fungicide appears to be less or no longer effective against a pathogen that it previously controlled or suppressed, contact a BASF representative, local extension specialist, or certified crop advisor to assist in determining the cause of reduced performance.

### Cleaning Spray Equipment

Clean spray equipment thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used prior to **Pristine® fungicide**.

### Application Instructions

Apply directed rates of **Pristine** as instructed by **Table 2. Pristine® fungicide Crop-specific Requirements**.

Ground application is recommended for thorough coverage. Aerial application can be made for those crops or in conditions where applications are not possible using ground equipment. **Pristine** can be applied through sprinkler irrigation equipment. Check equipment frequently for calibration. Under low-level disease conditions, use the minimum application rates; use maximum application rates and shortened spray schedules for severe or threatening disease conditions.

### Ground Application

Apply **Pristine** in sufficient water to ensure thorough coverage of foliage, bloom, and fruit. Thorough coverage is required for optimum disease control.

### Directed or Banded Sprays

The application rates on the **Pristine** label reflect the amount of product to be applied uniformly over an acre of ground on a broadcast basis.

In some crops, **Pristine** may be used as a directed or banded spray over the rows or plant beds with the alleys or row middles left unsprayed. For such uses, reduce the labeled **Pristine** rates in proportion to the area actually sprayed. This adjustment is necessary to avoid applying the product at use rates higher than permitted according to label directions.

Use the following formula to determine the broadcast equivalent rate for doing directed or banded sprays:

$$\begin{array}{r} \text{sprayed bed} \\ \text{width} \end{array} + \begin{array}{r} \text{unsprayed row} \\ \text{middles width} \end{array} = \text{total row width}$$

$$\frac{\begin{array}{r} \text{sprayed bed} \\ \text{width in inches} \end{array}}{\begin{array}{r} \text{total row} \\ \text{width in inches} \end{array}} \times \frac{\text{broadcast rate}}{\text{treated acre}} = \frac{\text{band rate}}{\text{field acre}}$$

**Example:** A directed spray application will be made to 45-inch plant beds that are separated by 15-inch unsprayed row middles.

$$\begin{array}{r} 45 \text{ inches} \\ \text{sprayed} \\ \text{bed width} \end{array} + \begin{array}{r} 15 \text{ inches} \\ \text{unsprayed} \\ \text{row middles} \end{array} = \begin{array}{r} 60 \text{ inches} \\ \text{total row width} \end{array}$$

The calculation to determine the appropriate equivalent rate of product to use for this situation based on a label broadcast rate recommendation of 12 ozs/acre follows:

$$\frac{\begin{array}{r} 45 \text{ inches} \\ \text{sprayed} \\ \text{bed width} \end{array}}{\begin{array}{r} 60 \text{ inches} \\ \text{total row width} \end{array}} \times \frac{\begin{array}{r} 12 \text{ ozs} \\ \text{Pristine} \\ \text{treated acre} \end{array}}{1} = \frac{\begin{array}{r} 9 \text{ ozs} \\ \text{Pristine} \\ \text{field acre} \end{array}}{1}$$

### Aerial Application

**For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).**

Aerial application can be made and thorough coverage is required to obtain optimum disease control. Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. Use no less than 5 gallons of spray solution per acre. For aerial applications to hops, tree and vine crops, use no less than 10 gallons of spray solution per acre. For all crops, thorough coverage is required for optimum disease control.

### Directions for Use Through Sprinkler Irrigation Systems

### Sprayer Preparation

Clean chemical tank and injector system thoroughly. Flush system with clean water.

### Application Instructions

Apply **Pristine** at rates and timings as described in this label.

### Use Precautions for Sprinkler Irrigation Applications

- This product can be applied through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Add this product to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop. **DO NOT** exceed 1/2 inch (13,577 gallons) per acre. In stationary or non-continuous moving systems, inject the product-water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Thorough coverage of

6/47

foliage is required for good control. Maintain good agitation during the entire application period.

- If you have questions about calibration, you should contact a state extension service specialist, equipment manufacturers or other experts.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- **DO NOT** connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

### Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the

flow of fluid back toward the injection pump.

4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

### Additives and General Tank Mixing Information

**Pristine® fungicide** can be tank mixed with most recommended fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives as specified in **Table 2. Pristine® fungicide Crop-specific Requirements**. See **Table 2. Pristine® fungicide Crop-specific Requirements** for exceptions.

Under some conditions, the use of additives or adjuvants may improve the performance of **Pristine**. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Pristine** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

### Compatibility Test and Mixing Order

If tank mixtures are used, adhere to restrictions due to rates, label recommendations and precautions on all labels.

### Compatibility Test for Tank Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of label rate per acre:

1. **Water** - For 100 gallons per acre spray volume, use 16 cups (1 gallon) of water. For other spray volumes, adjust

rates accordingly. Use only water from the intended source at the source temperature.

2. **Water-dispersible products** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions). Cap the jar and invert 10 cycles.
3. **Water-soluble products** - Cap the jar and invert 10 cycles.
4. **Emulsifiable concentrates** (oil concentrate or methylated seed oil when applicable). Cap the jar and invert 10 cycles.
5. **Water-soluble additives** - Cap the jar and invert 10 cycles.
6. Let the solution stand for 15 minutes.
7. **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

### Mixing Order

1. **Water** - Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (such as **Pristine® fungicide**, dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
6. **Water-soluble products**
7. **Emulsifiable concentrates** (such as oil concentrates when applicable)
8. **Water-soluble additives** (such as ammonium sulfate [AMS] or urea ammonium nitrate [UAN] when applicable)
9. **Remaining quantity of water**

Make sure that each component is thoroughly mixed and suspended before adding tank mix partners. Maintain constant agitation during application. See **Table 2. Pristine® fungicide Crop-specific Requirements** for more details.

### Restrictions and Limitations

- **DO NOT** exceed the maximum product rate (ozs/A) per year (season), the maximum product rate per application, or the total number of applications of **Pristine** per year (season) as stated in **Table 1. Pristine® fungicide Restrictions and Limitations Overview** and **Table 2. Pristine® fungicide Crop-specific Requirements**. Preharvest interval (PHI) restrictions are also included in these tables.
- **DO NOT** apply more than the maximum annual use rate

of ai/acre or ozs of product/acre for each specific crop from any combination of products containing pyraclostrobin or boscalid (e.g. **Pristine**, **Endura® fungicide**, **Cabrio® EG fungicide**, **Headline® fungicide**). To determine lbs of pyraclostrobin per acre, multiply ozs of product/acre by 0.008. To determine lbs of boscalid per acre, multiply ozs of product/acre by 0.0158.

- **Pristine** is not for use in greenhouse or transplant production.
- **Blueberry (highbush and lowbush) - DO NOT** apply **Pristine** to blueberries as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives or anything other than water. Mix **Pristine** with water only for applications to blueberries.
- **Grape - DO NOT** use on Concord or Noiret (NY73.0136.17) due to foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related grape varieties. Not all varieties have been thoroughly tested.
- **Aerial application in hops - DO NOT** make more than one (1) aerial application of **Pristine** per season and include a myclobutanil product as a tank mix.
- **For aerial application in New York State, DO NOT** apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

### Crop Rotation Restriction

Crops listed on the **Pristine**, **Cabrio EG**, **Endura** and **Headline** labels may be planted immediately following the last application.

All other crops can be planted 14 days after the last application.

**DO NOT** use on **cowpea, field pea, grain lupin, sugar beet, garden beet, turnip root or radish.**

9/47

Table 1. Pristine® fungicide Restrictions and Limitations Overview\*

Crop/Crop Group**	Minimum Time from Application to Harvest (PHI) (days)	Maximum Rate per Application (ozs product/A)	Maximum Number of Applications per Season***	Maximum Rate per Season (ozs product/A)
Alfalfa (including alfalfa grown for seed)	14	18	3	54
Belgium endive	19	1.6 (cold storage)	1	3.4
		1.8 (forcing)	1	
Berry subgroups	0	23	4	92
Bulb vegetables	7	18.5	6	111
Carrot	0	10.5	6	63
Celery	0	25	2	50
Citrus fruit	0	18.5	4	74
Cotton	30	25	2	50
Cucurbit vegetables	0	18.5	4	74
Dry beans (except soybean)	21	25	2	50
Fruiting vegetables	0	9.7	6	58.2
Tomato***		25	2	69
Globe artichoke	0	23	3	69
Grape***	14	23	3	69
Hops****	14	28	3	84
Leafy greens (except <i>Brassica</i> and head lettuce and leaf lettuce)	14	25	2	50
Leafy petioles (except <i>Brassica</i> )	0	25	2	50
Low growing berry subgroup (except cranberry and strawberry)	0	23	5	115
Oilseed crops***	21	24.5	2	49
Peanut	14	28	3	84
Persimmon	0	23	3	69
Pome fruit	0	18.5	4	74

(continued)

10/47

**Table 1. Pristine® fungicide Restrictions and Limitations Overview\*** (continued)

Crop/Crop Group**	Minimum Time from Application to Harvest (PHI) (days)	Maximum Rate per Application (ozs product/A)	Maximum Number of Applications per Season***	Maximum Rate per Season (ozs product/A)
Small fruit, vine climbing subgroup*** (except fuzzy kiwifruit and grape)	14	23	5	69
Soybean	21	16	2	32
Spinach	14	25	2	50
Stone fruit	0	14.5	5	72.5
Strawberry	0	23	5	115
Tree nut	14 (for almond - 25 days)	14.5	4	58
Tropical fruit	0	18.5	2	37
Turnip greens	14	25	2	50

\* See **Table 2. Pristine® fungicide Crop-specific Requirements** for complete directions and exceptions, including restrictions and recommendations regarding crop sensitivity as well as tank mixtures.

\*\* For a complete list of crops labeled within a group, see **Table 2. Pristine® fungicide Crop-specific Requirements**.

\*\*\* At maximum use rate only, except for grape; oilseed crops; small fruit, vine climbing; and tomato.

\*\*\*\* For additional ground and/or aerial application restrictions and limitations, see **Table 2. Pristine® fungicide Crop-specific Requirements, Hops**.

Aerial application is permitted for all labeled crop uses. **For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).**

**Crop-specific Requirements**

**Table 2. Pristine® fungicide Crop-specific Requirements**

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Alfalfa (including alfalfa grown for seed)</b>	Anthracnose ( <i>Colletotrichum trifolii</i> )	12 to 18	2 per cutting or 3 total per growing season	54	14
	Common leaf spot ( <i>Pseudopeziza medicaginis</i> )				
	Downy mildew ( <i>Peronospora trifoliorum</i> )				
	Leaf spot ( <i>Leptosphaerulina briosiani</i> )				
	Powdery mildew ( <i>Erysiphe pisi</i> )				
	Rhizoctonia blight/Black patch ( <i>Rhizoctonia</i> spp.)				
	Rust ( <i>Uromyces</i> spp.)				
	Spring black stem and Leaf spot ( <i>Phoma medicaginis</i> )				
	Stagonospora leaf spot ( <i>Stagonospora melloti</i> )				
	Stemphyllium leaf spot ( <i>Stemphyllium</i> spp.)				
	Summer black stem and Leaf spot ( <i>Cercospora medicaginis</i> )	14 to 18			
	Yellow leaf blotch ( <i>Leptotrichila medicaginis</i> )				
	White mold/Sclerotinia crown and Stem rot ( <i>Sclerotinia sclerotiorum</i> , <i>S. trifoliorum</i> )				
	<b>Suppression Only:</b> Southern blight ( <i>Sclerotium rolfsii</i> )				

(continued)

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)**Alfalfa** (continued)

**Application Directions.** Begin **Pristine** applications when conditions favorable for disease are expected, but prior to onset of disease development. For stand establishment of fall-seeded alfalfa, begin applications in fall through early winter prior to first snowfall or extended cool, wet conditions. For seed pod protection, begin applications at 10% to 30% bloom.

Using higher rates may improve disease control performance as the crop canopy volume and density increases. Disease control can also be improved when application equipment and spray volume is adjusted to achieve thorough canopy penetration and coverage.

Repeat application on a 14 to 21 day interval if conditions are favorable for disease development. **DO NOT** make more than two (2) **Pristine** applications per cutting or more than three (3) **Pristine** applications per season.

Use the higher rate and shorter interval when disease pressure is high.

Under some conditions, additives or adjuvants may improve the performance of **Pristine**.

No livestock feeding restrictions.

**Resistance Management.** To limit development of resistance, **DO NOT** make more than two (2) sequential **Pristine** applications per cutting or three (3) **Pristine** applications per season. Alternate to a labeled **non-Group 7** or **non-Group 11** fungicide with different mode of action following two (2) sequential **Pristine** applications.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application	Maximum Number of Applications per Season	Maximum Product Rate per Crop (ozs product per 1000 lbs roots)	Minimum Time from Application to Harvest (PHI) (days)
Belgium endive*	Root and crown rot ( <i>Sclerotinia sclerotiorum</i> )	<b>Prior to cold storage:</b> 0.8 to 1.6 oz per 1000 lbs roots	1	3.4	19
		<b>Prior to forcing:</b> 0.9 to 1.8 oz per 70 square feet of forcing tray	1		

**Application Directions. Dosage and frequency/timing of applications.** Make one application to the roots when brought into cold storage prior to forcing. Apply again at the beginning of forcing after the roots have been packed in forcing trays.

**Prior to Cold Storage.** Make one application as a spray to the roots as they move along a conveyor belt used to bring roots from field transportation into cold storage bins. Apply 0.8 to 1.6 ozs **Pristine** in 3.0 to 3.5 gals of water per 1000 lbs roots.

**Prior to Forcing.** Make one application as a spray to the roots at the beginning of forcing, after they have been packed into forcing trays. Apply at the rate of 0.9 to 1.8 ozs of **Pristine** in approximately 100 fl ozs of water per 70 square feet of forcing tray. Approximately 1000 lbs of roots will fill 70 square feet of forcing tray.

**Restrictions. DO NOT** apply after the beginning of forcing.

\* For use in California only.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Berry subgroups*</b>	Alternaria leaf spot and fruit rot ( <i>Alternaria</i> spp.)	.18.5 to 23	4	92	0
<b>Bushberry subgroup</b>					
Aronia berry	Anthracnose				
Black currant	( <i>Colletotrichum</i> spp.,				
Blueberry**	<i>Elsinoe</i> spp.)				
(highbush and lowbush)	Botrytis gray mold				
Buffalo currant	( <i>Botrytis cinerea</i> )				
Chilean guava	Leaf spot and blotch				
Currant	( <i>Mycosphaerella</i> spp.,				
Elderberry	<i>Septoria</i> spp.)				
European barberry	Monilinia blight and mummy				
Gooseberry	berry				
Highbush cranberry	( <i>Monilinia</i> spp.)				
Honeysuckle, edible	Phomopsis leaf spot,				
Huckleberry	twig blight, and fruit rot				
Jostaberry	( <i>Phomopsis</i> spp.)				
Juneberry	Powdery mildew				
Lingonberry	( <i>Sphaerotheca</i> spp.,				
Native currant	<i>Microsphaera</i> spp.,				
Red currant	<i>Oidium</i> spp.)				
Salal	Spur blight				
Sea buckthorn	( <i>Didymella</i> spp.,				
<b>Caneberry subgroup</b>	<i>Phoma</i> spp.)				
Blackberry	<b>Suppression Only:</b>				
(all varieties)	Rust				
Loganberry	( <i>Pucciniastrum</i> spp.,				
Raspberry	<i>Arthuriomyces</i> spp.,				
(black and red)	<i>Phragmidium</i> spp.,				
Wild raspberry	<i>Kuehneola</i> spp.)				

**Application Directions.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

\* **For the berries listed in the berry subgroups (except blueberry) in this table,** it is impossible for BASF to test all Berries Group crops for sensitivity to **Pristine** under all environments and all potential product mixture combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Pristine** use, particularly in tank mixes and/or adjuvant combinations on berry crops. To reduce the risk of berry crop injury, BASF recommends testing **Pristine** or **Pristine** tank mixtures on a small portion of the crop before broad scale use. To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Pristine** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

\*\* **Blueberry (highbush and lowbush)** is not registered for use in California. For all other states, **DO NOT** apply **Pristine** to blueberries as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives or anything other than water. Mix **Pristine** with water only for applications to blueberries.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Bulb vegetables group</b> Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem	Botrytis leaf blight ( <i>Botrytis</i> spp.) Botrytis neck rot* ( <i>Botrytis</i> spp.)	14.5 to 18.5	6	111	7
	Purple blotch and leaf blight ( <i>Alternaria porri</i> )	10.5 to 18.5			
	<b>Suppression Only:</b> Downy mildew ( <i>Peronospora destructor</i> )	18.5			
Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties and/or hybrids of these					

**Application Directions.** For control of neck rot, purple blotch and leaf blight, begin applications of **Pristine** prior to onset of disease development and continue on a 14 day interval. If application intervals shorter than 14 days are needed, rotate to another fungicide with a different mode of action.

Use the higher rate when disease pressure is high.

Applications made to control purple blotch, leaf blight and stalk rot will also suppress downy mildew. If downy mildew occurs during a **Pristine** application for these diseases, immediately follow the **Pristine** application with a downy mildew fungicide with a different mode of action.

**For downy mildew,** rotate each application of **Pristine** with an application of a labeled fungicide with a different mode of action.

No restriction on livestock grazing or feeding.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than six (6) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

\* Not registered for use in California.

16/47

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Carrot	Alternaria leaf spot ( <i>Alternaria</i> spp.)  Cercospora leaf spot ( <i>Cercospora</i> spp.)  Powdery mildew ( <i>Erysiphe</i> spp.)	8 to 10.5	6	63	0
	<b>Suppression Only:</b>  Southern root rot ( <i>Sclerotium rolfsii</i> )				

**Application Directions.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7 to 14 day interval.

Use the higher rate and the shorter interval when disease pressure is high.

No restriction on livestock grazing or feeding for carrot culls.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than six (6) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per crop growing season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Celery*</b> <b>Celery (Chinese)*</b>	Alternaria leaf spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum</i> spp.) Ascochyta leaf spot ( <i>Ascochyta</i> spp.) Cercospora leaf spot ( <i>Cercospora</i> spp.) Downy mildew ( <i>Peronospora</i> spp., <i>Bremia</i> spp.) Phoma ( <i>Phoma</i> spp.) Rust ( <i>Puccinia</i> spp.) Powdery mildew ( <i>Erysiphe</i> spp.) Septoria leaf spot ( <i>Septoria</i> spp.) White rust ( <i>Albugo</i> spp.)	10 to 15	2	50	0
	Botrytis rot ( <i>Botrytis</i> spp.) Sclerotinia rot and blight ( <i>Sclerotinia</i> spp.)	25			

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7 day interval.

Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

\* Not registered for use in California.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Citrus fruit group</b> Australian desert lime Australian finger lime Australian round lime Brown River finger lime Calamondin Chironja Citron Citrus hybrids Grapefruit Japanese summer grapefruit Kumquat Lemon Lime Mediterranean mandarin Mount white lime New Guinea wild lime Orange, sour Orange, sweet Pummelo Russell River lime Satsuma mandarin Sweet lime Tachibana orange	Alternaria brown spot <i>(Alternaria alternata, Alternaria spp.)</i>  Citrus black spot* <i>(Guignardia citricarpa)</i>  Greasy spot <i>(Mycosphaerella citri)</i>  Melanose <i>(Diaporthe citri)</i>  Scab <i>(Elsinoe fauceitii)</i>	16 to 18.5	4	74	0
Tahiti lime Tangelo Tangerine (mandarin) Tangor Trifolate orange Uniq fruit Cultivars, varieties and/or hybrids of these					

**Application Directions.** Apply **Pristine** in a regularly scheduled protective fungicide program. Begin **Pristine** applications prior to infection and continue on a 10 to 21 day interval.

Use the higher rate and shorter interval when disease pressure is high.

Disease control from **Pristine** depends on disease pressure and various cultural practices that influence rind maturation and disease susceptibility. Improved disease performance may result when **Pristine** is used in a crop management program that minimizes rind overmaturity and rind damage.

No livestock feeding restrictions.

**Resistance Management:** To limit development of resistance, **DO NOT** make more than four (4) **Pristine** applications per season.

**DO NOT** make more than two (2) sequential **Pristine** applications before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide with different modes of action.

\* Not registered for use in California.

19/  
47

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Cotton</b>	Alternaria leaf spot ( <i>Alternaria</i> spp.)  Anthracnose, boll rot ( <i>Glomerella</i> spp.)  Ascochyta blight, boll rot ( <i>Ascochyta</i> spp.)  Cercospora leaf spot ( <i>Cercospora</i> spp.)  Diplodia boll rot ( <i>Diplodia</i> spp.)  Hard lock, boll rot ( <i>Fusarium</i> spp.)	12.5 to 25	2	50	30
<p><b>Application Directions.</b> Begin applications of <b>Pristine</b> prior to the onset of disease development and continue on a 7 to 14 day interval.</p> <p>Use the higher rate and the shorter interval when disease pressure is high.</p> <p>Feed containing commodities from cotton production and processing can be fed to livestock.</p> <p><b>Resistance Management.</b> To limit the potential for development of resistance, <b>DO NOT</b> make more than one (1) application of <b>Pristine</b> before alternating to a labeled fungicide with a different mode of action for at least one (1) application.</p>					

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Cucurbit vegetables group</b>  <b>Includes all types and hybrids of:</b>                      Chayote                      Chinese waxgourd                      Citron melon                      Cucumber                      Gherkin                      Pumpkin                      Watermelon</p> <p><b>Edible Gourd</b>                      Chinese okra                      Cucuzza                      Hyotan</p> <p><b>Momordica spp.</b>                      Balsam apple                      Balsam pear                      Bitter melon                      Chinese cucumber</p> <p><b>Muskmelon</b>                      Cantaloupe                      Casaba                      Grenshaw-melon</p>	<p>Downy mildew (<i>Pseudoperonospora cubensis</i>)                      Alternaria blight (<i>Alternaria cucumerina</i>)                      Cercospora leaf spot (<i>Cercospora citrulina</i>)                      Gummy stem blight (<i>Didymella bryoniae</i>)                      Powdery mildew (<i>Sphaerotheca fuliginea</i>, <i>Erysiphe cichoracearum</i>)</p>	<p>12.5 to 18.5</p>	<p>4</p>	<p>74</p>	<p>0</p>
<p>Golden pershaw melon                      Honeydew melon                      Honey balls                      Mango melon                      Persian melon                      Pineapple melon                      Santa Claus melon                      Snake melon</p> <p><b>Summer Squash</b>                      Crookneck squash                      Scallop squash                      Straightneck squash                      Vegetable marrow                      Zucchini</p> <p><b>Winter Squash</b>                      Acorn squash                      Butternut squash                      Calabaza                      Hubbard squash                      Spaghetti squash</p>	<p>Anthracoise (<i>Colletotrichum orbiculare</i>)</p>	<p>18.5</p>			

(continued)

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

**Cucurbit vegetables group** (continued)

**Application Directions.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7 to 14 day interval.

Use the higher rate and the shorter interval when disease pressure is high.

Use the highest labeled rate for anthracnose.

**Tank Mixes with Adjuvants and Other Products.** BASF evaluations indicate that tank mixes of additives, adjuvants, and/or other products with **Pristine** may result in injury. This is particularly true for muskmelon crops such as cantaloupe and honeydew. Users need to be aware of this, proceed with caution, and test for crop safety when tank mixing, as stated below.

Applications of additives, adjuvants, and/or other products that increase penetration may cause injury when mixed with **Pristine**. Injury potential from these kinds of tank mixes may decrease with lower rates of the tank mix partner. Users are advised to test for crop safety, as stated below.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives, adjuvants, and/or other products. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Pristine** with other products.

To minimize the likelihood of crop injury, BASF recommends testing **Pristine** in combination with additives, adjuvants, and/or other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

**DO NOT** tank mix **Pristine** with malathion, **Kelthane® agricultural miticide**, **Thiodan® insecticide**, **Phaser® insecticide**, **Lannate® insecticide**, **Lorsban® insecticide**, **M-Pede® insecticide/fungicide**, or **Botran® fungicide** as crop injury may result.

**Resistance Management.** To limit the potential of development of resistance, **DO NOT** make more than four (4) applications of **Pristine** per season.

~~**DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.~~



**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Fruiting vegetables group</b> African eggplant Bush tomato Bell pepper Cocona Currant tomato Eggplant Garden huckleberry Goji berry Groundcherry Martynia Naranjilla Okra Pea eggplant Pepino Pepper (all varieties) Nonbell pepper Roselle Scarlet eggplant Sunberry Tomatillo Tree tomato	Black mold ( <i>Alternaria alternata</i> ) Early blight ( <i>Alternaria solani</i> )	9.7 or 9.7 ozs per 100 gal of spray volume (dilute)*	6	58.2	0
Cultivars, varieties and/or hybrids of these					
<b>Tomato</b>	Anthracnose ( <i>Colletotrichum</i> spp.) Black mold ( <i>Alternaria alternata</i> ) Botrytis gray mold ( <i>Botrytis cinerea</i> ) Early blight ( <i>Alternaria solani</i> ) Late blight ( <i>Phytophthora infestans</i> ) Powdery mildew ( <i>Leveillula taurica</i> ) Septoria leaf spot ( <i>Septoria lycopersici</i> ) Target spot ( <i>Corynespora cassiicola</i> )	12.5 to 25	5 at 12.5 ozs/A 2 at 25 ozs/A	69	

(continued)

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)**Fruiting vegetables group** (continued)

**Application Directions.** Begin **Pristine** applications prior to disease development and continue on a 7 to 14 day interval for anthracnose, black mold, botrytis gray mold, early blight, powdery mildew, Septoria leaf spot, and target spot. For control of late blight, begin applications prior to disease development; then follow each **Pristine** application with a labeled fungicide with a different mode of action 5 to 7 days later.

Use the higher rate and shorter interval when disease pressure is high.

\* For applications based on dilute volume, spray plants to runoff. Apply a minimum of 20 gallons of spray volume per acre, and increase the spray volume as the plants grow during the season. Spray proportional volume to the amount of plant tissue to be covered such that 100 gallons of spray per acre is used on mature plants.

**Use of Adjuvants.** Additive or adjuvant use may improve the performance of **Pristine** on fruiting vegetables. However, BASF evaluations also indicate that under some conditions (particularly high temperatures and/or high additive rates), **Pristine** application in combination with certain rates of silicone based or oil-containing (petroleum or crop) additives or adjuvants can cause injury.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Pristine** with other products.

To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Pristine** spray solution. BASF cannot be held responsible for crop injury, reduced disease control or incompatibility because of additives, adjuvants or other products used in combination with **Pristine** (see **Conditions of Sale and Warranty**).

To minimize the likelihood of crop injury, BASF recommends testing **Pristine** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

**Resistance Management.** To limit development of resistance, **DO NOT** apply more than 1.2 lbs ai pyraclostrobin per acre per crop growing season. **DO NOT** make more than two (2) sequential **Pristine** applications before alternating to a-labeled ~~non-Group 7 or non-Group 11~~ fungicide with a different mode of action.

For additional resistance management information, see **Resistance Management** in the **Product Information** section.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Globe artichoke</b>	Bud rot ( <i>Botrytis cinerea</i> )  Ramularia leaf spot ( <i>Ramularia</i> spp.)	18.5 to 23.0	3	69	0

**Application Directions. Dosage and frequency/timing of applications.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7 to 14 day interval. For artichoke bud rot, begin applications at the initiation of the bud protection phase when approximately 25% of the plants have bolted. Use the shorter interval and/or the higher rate when disease pressure is high.

**DO NOT** apply **Pristine** to artichokes as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives or anything other than water. Mix **Pristine** with water only for applications to artichokes.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** exceed the specified number of applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season. Adhere to the label instructions regarding the consecutive use of **Pristine** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

26/47

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Grape (except Concord or Noiret (NY73.0136.17) due to foliar injury.</b> It is possible that foliar injury could occur on related grape varieties. See comments in the <b>Application Directions</b> below for more information.	Angular leaf spot ( <i>Mycosphaerella angulata</i> ) Anthracnose ( <i>Elsinoe ampelina</i> ) Black rot ( <i>Guignardia bidwellii</i> ) Downy mildew ( <i>Plasmopara viticola</i> ) Leaf blight ( <i>Pseudocercospora vitis</i> ) Phomopsis cane and leaf spot ( <i>Phomopsis viticola</i> ) Powdery mildew ( <i>Uncinula necator</i> ) Ripe rot ( <i>Colletotrichum gloeosporioides</i> ) <b>Aids in Control Only:</b> Summer bunch rot (Sour rot) ( <i>Cladosporium</i> spp. and <i>Aspergillus</i> spp.) <b>Suppression Only:</b>	8 to 12.5	5	69	14
	Botrytis gray mold ( <i>Botrytis cinerea</i> ) Botrytis gray mold ( <i>Botrytis cinerea</i> )				

**Application Directions.** For powdery mildew control, begin applications of **Pristine** as of bud break prior to onset of disease, using 8 ozs per acre on a 10 to 14 day interval. Use 10 to 12.5 ozs per acre on a 14 to 21 day interval.

**For black rot and downy mildew control,** begin applications of **Pristine** as of pre-bloom prior to onset of disease and continue applications on a 10 to 14 day interval.

**For all other diseases listed except for Botrytis gray mold,** begin applications of **Pristine** prior to onset of disease and continue applications on a 10 to 14 day interval. **Pristine** applied at rates of 8 to 12.5 ozs per acre for control of the listed diseases will also suppress Botrytis gray mold.

**For control of Botrytis gray mold,** apply 18.5 to 23 ozs per acre of **Pristine** prior to onset of disease development when conditions favor disease development during early bloom, bunch pre-closure and veraison up to 14 days before harvest. Use the higher rate and the shorter interval when disease pressure is high.

**DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours except when performing cane tying, cane turning or cane girdling. The REI is 5 days for treated grapes when conducting cane tying, cane turning or cane girdling.**

**DO NOT use on Concord or Noiret (NY73.0136.17) due to foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related grape varieties. Not all varieties have been thoroughly tested.** Consult a BASF representative for more information concerning these sensitive grapes.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application	Maximum Number of Ground Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Hops (Ground Application)</b>	Powdery mildew ( <i>Erysiphe cichoracearum</i> , <i>Sphaerotheca</i> spp.)  Downy mildew ( <i>Pseudoperonospora humuli</i> )	14 ozs per 100 gallons of dilute spray  <b>DO NOT</b> use more than 28 ozs per acre	3  (2 if one aerial application is made)	84  (70 ozs/A if one aerial application is made)	14

**Application Directions.** Begin applications of **Pristine** prior to disease development and continue on a 10 to 21 day interval.

Use the shorter interval when disease pressure is high.

Application rates are based on 100 gallons of dilute spray applied to runoff. Adjust water volume to maintain thorough coverage. Use 25 to 50 gallons of dilute spray per acre prior to trellising and 100 to 200 gallons of dilute spray per acre thereafter. **DO NOT** use more than 200 gallons per acre of this mixture. If additional spray volume is needed for thorough coverage, use 28 ozs of **Pristine** per acre in the required spray volume.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than three (3) applications of **Pristine** per season (counting both ground and aerial applications). **DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Restrictions.** **DO NOT** use more than 200 gallons per acre of this mixture. If additional spray volume is needed for thorough coverage, use 28 ozs of **Pristine** per acre in the required spray volume.

Crop	Target Disease	Aerial Product Use Rate per Application	Maximum Number of Aerial Applications per Season	Aerial Application Timing Growth Stage	Minimum Time from Aerial Application to Harvest (PHI) (days)
<b>Hops (Aerial Application)</b>	Powdery mildew ( <i>Erysiphe cichoracearum</i> , <i>Sphaerotheca humuli</i> , <i>Sphaerotheca macularis</i> , <i>Sphaerotheca</i> spp.)	14 ozs/A as a tank mix with a myclobutanil fungicide product (see myclobutanil rate following)	1	Wire to 14 days preharvest	14

**Application Directions.** Aerial application may result in reduced control due to lack of canopy penetration and coverage. Aerial application should only be used in situations when ground application is not possible.

Apply a preventive spray of **Pristine** at 14 ozs as a tank mix with a myclobutanil fungicide product at rates **equivalent to 0.15 lb per acre of active ingredient** (including but not limited to: **Rally® 40W fungicide** at 6 ozs product per acre or **Sonoma® 40WSP fungicide** at 6 ozs product per acre) for resistance management.

Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. Use a minimum of 10 gallons of water per acre when applying by air. Thorough coverage is essential.

Because complete coverage is important for effective disease control, **aerial application at low volumes may result in reduced control due to lack of canopy penetration and coverage.**

Mixing **Pristine** with surfactants or foliar fertilizers is not recommended when applying by air. Similarly, adjuvants that enhance pesticide penetration may cause phytotoxicity when used with **Pristine** applied by air.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) aerial application of **Pristine** per season and include a myclobutanil product as a tank mix as described.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Leafy greens* (except Brassica and head lettuce and leaf lettuce)</b> Amaranth Arugula Chervil Chrysanthemum (edible-leaved and garland) Corn salad Cress (garden and upland) Dandelion Dock Endive Orach Parsley Purslane (garden and winter) Radicchio (red chicory)	Alternaria leaf spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum</i> spp.) Ascochyta leaf spot ( <i>Ascochyta</i> spp.) Cercospora leaf spot ( <i>Cercospora</i> spp.) Downy mildew ( <i>Peronospora</i> spp.) Phoma ( <i>Phoma</i> spp.) Powdery mildew ( <i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.) Rust ( <i>Puccinia</i> spp.) Septoria leaf spot ( <i>Septoria</i> spp.)	10 to 15	2	50	14
	White rust ( <i>Albugo</i> spp.)				
	Botrytis rot ( <i>Botrytis</i> spp.)	15 to 25			
	Sclerotinia rot and blight ( <i>Sclerotinia</i> spp.)				

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7 day interval.

Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

\* Not registered for use in California except on radicchio for control of Sclerotinia rot and blight when applied at a rate of 25 ozs/A.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Leafy petioles (except Brassica)</b> Cardoon Celtnce Fennel (Florence) Rhubarb Swiss chard	Alternaria leaf spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum</i> spp.) Ascochyta leaf spot ( <i>Ascochyta</i> spp.) Cercospora leaf spot ( <i>Cercospora</i> spp.) Downy mildew ( <i>Peronospora</i> spp., <i>Bremia</i> spp.) Phoma ( <i>Phoma</i> spp.) Powdery mildew ( <i>Erysiphe</i> spp.) Rust ( <i>Puccinia</i> spp.) Septoria leaf spot ( <i>Septoria</i> spp.) White rust ( <i>Albugo</i> -spp.)	10 to 15	2	50	0
	Botrytis rot ( <i>Botrytis</i> spp.) Sclerotinia rot and blight ( <i>Sclerotinia</i> spp.)	15 to 25			

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7 day interval.

Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Low growing berry subgroup (except cranberry and strawberry)</b> Bearberry Bilberry Cloudberry Muntries Partridgeberry	Anthracnose ( <i>Colletotrichum</i> spp.) Botrytis gray mold ( <i>Botrytis cinerea</i> ) Leaf spot ( <i>Mycosphaerella fragariae</i> ) Powdery mildew ( <i>Sphaerotheca macularis</i> )	18.5 to 23	5	115	0

**Application Directions.** Begin applications of **Pristine** no later than 10% bloom, or prior to disease development, and continue on a 7 to 14 day interval.

Use the higher rate and the shorter interval when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oilseed crops Flax seed	PasmO ( <i>Septoria linicola</i> )	16.7	2	33.4	21
Rapeseed (canola varieties only)	Blackleg ( <i>Leptosphaeria maculans</i> )  Blackspot ( <i>Alternaria</i> spp.)  Sclerotinia rot and blight ( <i>Sclerotinia</i> spp.)				
Sunflower	Alternaria leaf spot ( <i>Alternaria</i> spp.)  Cercospora leaf spot ( <i>Cercospora helianthi</i> )  Downy mildew ( <i>Plasmopara halstedii</i> )  Powdery mildew ( <i>Erysiphe cichoracearum</i> )  Rust ( <i>Puccinia helianthi</i> , <i>Uromyces</i> spp.)  Sclerotinia rot and blight ( <i>Sclerotinia</i> spp.)	24.5	2	49	21
	Septoria leaf spot ( <i>Septoria</i> spp.)  White rust ( <i>Albugo tragopogonis</i> )				

(continued)

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Oilseed crops</b> (continued) Borage Calendula Castor oil plant Chinese tallowtree Crambe Cuphea Echium Euphorbia Evening primrose Gold of pleasure Hare's ear mustard Jojoba Lesquerella Lunaria Meadowfoam Milkweed Mustard seed Niger seed Oil radish Poppy seed Rose hip Safflower Sesame Stokes aster Sweet rocket Tallowwood Tea oil plant Vernonia	Alternaria spp. Septoria spp.	16.7	2	33.4	21

**Application Directions.** For optimal disease control, begin applications of **Pristine** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

**Rapeseed. For control of blackleg,** apply **Pristine** at 2- to 4-leaf stage. **For optimal control of blackspot,** apply **Pristine** at early pod development. A second application 7 to 10 days later may be made if disease persists or if weather conditions are favorable for disease development.

**Flax seed.** Apply **Pristine** at mid-flowering (7 to 10 days after flower initiation). Make a second application 7 to 10 days later if disease persists or if weather conditions are favorable for disease development.

**Pristine** may be used with adjuvants.

No livestock feeding restrictions.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) applications per season. **DO NOT** make more than two (2) applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Peanut*	Early leaf spot ( <i>Cercospora arachidicola</i> )  Late leaf spot ( <i>Cercosporidium personatum</i> )  Pepperspot ( <i>Leptosphaerulina crassiasca</i> )  Rust ( <i>Puccinia arachidis</i> )  Web blotch ( <i>Phoma arachidicola</i> )	12.5 to 18.5	3	84	14
	Rhizoctonia limb rot, peg rot, and pod rot ( <i>Rhizoctonia solani</i> )  Sclerotium rot, Southern stem rot, Southern blight, and white mold ( <i>Sclerotium rolfsii</i> )  Sclerotinia blight ( <i>Sclerotinia minor</i> )	18.5 to 28			

**Application Directions.** For control of pepperspot, rust, web blotch, early and late leaf spot, begin applications of **Pristine** prior to onset of disease development and continue on a 14 day interval.

For control of **Rhizoctonia** and **Sclerotium** rot, begin applications of **Pristine** prior to onset of disease development and continue on a 14 day interval.

For control of **Sclerotinia** blight, begin applications of **Pristine** prior to onset of disease development or 45 to 60 days after planting. Make a second application 14 to 21 days later.

Use the higher rate and/or shorter spray interval when disease pressure is high or in fields with a history of disease.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than three (3) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Restrictions:** Use of **Pristine** with silicone-based adjuvants may cause crop injury.

**DO NOT** feed treated peanut hay to livestock.

**DO NOT** graze livestock or harvest for forage use.

\* Not registered for use in California.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Persimmon	Cercospora leaf spot ( <i>Cercospora</i> spp.)	18.5 to 23.0	3	69	0

**Application Directions. Dosage and frequency/timing of applications.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** exceed the specified number of applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season. Adhere to the label instructions regarding the consecutive use of **Pristine** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Pome fruit group</b>	Alternaria blotch ( <i>Alternaria mali</i> )	14.5 to 18.5	4	74	0
Apple					
Azarole	Apple scab ( <i>Venturia inaequalis</i> )				
Crabapple					
Loquat	Bitter rot ( <i>Colletotrichum</i> spp.)				
Mayhaw					
Medlar					
Pear	Black rot/Frogeye leaf spot ( <i>Botryosphaeria obtusa</i> )				
Pear, Asian					
Pear, Oriental	Blue mold* ( <i>Penicillium</i> spp.)				
Quince					
Quince, Chinese	Brooks spot ( <i>Mycosphaerella pomi</i> )				
Quince, Japanese					
Tejocote					
Cultivars, varieties and/or hybrids of these	Flyspeck ( <i>Zygophiala jamaicensis</i> )				
	Gray mold* ( <i>Botrytis</i> spp.)				
	Pear scab ( <i>Venturia pirina</i> )				
	Powdery mildew ( <i>Podosphaera leucotricha</i> )				
	Sooty blotch (disease complex)				
	White rot ( <i>Botryosphaeria dothidea</i> )				
	<b>Suppression Only:</b>				
	Cedar apple rust ( <i>Gymnosporangium juniperi-virginianae</i> )				
	Quince rust ( <i>Gymnosporangium clavipes</i> )				

**Application Directions for scab, powdery mildew, frogeye leaf spot and rust.** Begin applications of **Pristine** prior to disease development and continue on a 7 to 10 day interval.

Use the higher rate and shorter interval when disease pressure is high.

**Application Directions for blue mold, gray mold, sooty blotch, flyspeck, white rot, black rot, bitter rot and Alternaria blotch.** Begin applications of **Pristine** prior to disease development and continue on a 7 to 14 day interval.

Use the higher rate and shorter interval when disease pressure is high.

**For pears, DO NOT** use **Pristine** with a horticultural mineral oil as crop response to foliage and/or fruit can occur under certain conditions.

No restriction on livestock grazing or feeding for pome fruits feed items.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

\* Not registered for use in California.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Small fruit, vine climbing subgroup (except fuzzy kiwifruit and grape)</b> Amur river grape Gooseberry Kiwifruit, hardy Maypop Schisandra berry	Angular leaf spot ( <i>Mycosphaerella angulata</i> )	8 to 12.5	5	69	14
	Anthracnose ( <i>Elsinoe ampelina</i> )				
	Black rot ( <i>Guignardia bidwellii</i> )				
	Downy mildew ( <i>Plasmopara viticola</i> )				
	Leaf blight ( <i>Pseudocercospora vitis</i> )				
	Phomopsis cane and leaf spot ( <i>Phomopsis viticola</i> )				
	Powdery mildew ( <i>Uncinula necator</i> )				
	Ripe rot ( <i>Colletotrichum gloeosporioides</i> )				
	<b><u>Aids in Control Only:</u></b> Summer bunch rot (Sour rot) ( <i>Cladosporium</i> spp. and <i>Aspergillus</i> spp.)				
	<b><u>Suppression Only:</u></b> Botrytis gray mold ( <i>Botrytis cinerea</i> )				
	Botrytis gray mold ( <i>Botrytis cinerea</i> )	18.5 to 23	3		

**Application Directions. For powdery mildew control,** begin applications of **Pristine** as of bud break prior to onset of disease, using 8 ozs per acre on a 10 to 14 day interval. Use 10 to 12.5 ozs per acre on a 14 to 21 day interval.

**For black rot and downy mildew control,** begin applications of **Pristine** as of pre-bloom prior to onset of disease and continue applications on a 10 to 14 day interval.

**For all other diseases listed except for Botrytis gray mold,** begin applications of **Pristine** prior to onset of disease and continue applications on a 10 to 14 day interval. **Pristine** applied at rates of 8 to 12.5 ozs per acre for control of the listed diseases will also suppress Botrytis gray mold.

**For control of Botrytis gray mold,** apply 18.5 to 23 ozs per acre of **Pristine** prior to onset of disease development when conditions favor disease development during early bloom, bunch pre-closure and veraison up to 14 days before harvest.

Use the higher rate and the shorter interval when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Soybean*	Alternaria leaf spot ( <i>Alternaria</i> spp.)	8 to 16	2	32	21
	Anthracnose ( <i>Colletotrichum truncatum</i> )				
	Brown spot ( <i>Septoria glycines</i> )				
	Cercospora blight ( <i>Cercospora kikuchii</i> )				
	Frogeye leaf spot ( <i>Cercospora sojina</i> )				
	Pod and stem blight ( <i>Diaporthe phaseolorum</i> )				
	Rhizoctonia aerial blight ( <i>Rhizoctonia solani</i> )				
	Asian soybean rust ( <i>Phakopsora pachyrhizi</i> )	12.5 to 16			
Southern blight ( <i>Sclerotium rolfsii</i> )	16				
White mold ( <i>Sclerotinia sclerotiorum</i> )					

**Application Directions.** For optimal disease control, apply **Pristine** at early flowering (R1 to R3 growth stage) or prior to disease development, whichever is earlier. Make a second application 7 to 21 days later if monitoring shows disease development or if conditions are conducive for disease infection.

Use the higher labeled rate and shorter interval when disease pressure is high.

**Pristine** may be applied with adjuvants.

Soybean forage may be fed no sooner than 14 days after last application. Soybean hay may be fed no sooner than 21 days after last application.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

\* Not registered for use in California.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Spinach</b> <b>Spinach (New Zealand and vine)</b>	Alternaria leaf spot ( <i>Alternaria</i> spp.)	10 to 15	2	50	14
	Anthracnose ( <i>Colletotrichum</i> spp.)				
	Ascochyta leaf spot ( <i>Ascochyta</i> spp.)				
	Cercospora leaf spot ( <i>Cercospora</i> spp.)				
	Downy mildew ( <i>Peronospora</i> spp.)				
	Phoma ( <i>Phoma</i> spp.)				
	Powdery mildew ( <i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.)				
	Rust ( <i>Puccinia</i> spp.)				
	Septoria leaf spot ( <i>Septoria</i> spp.)				
	White rust ( <i>Albugo</i> spp.)				
	Botrytis rot ( <i>Botrytis</i> spp.)	25			
	Sclerotinia rot and blight ( <i>Sclerotinia</i> spp.)				

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7 day interval.

Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Stone fruit group</b> Apricot Cherry (sweet and tart) Nectarine Peach Plum (all varieties) Plumcot Prune	Alternaria leaf spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum</i> spp.) Blossom blight ( <i>Monilinia</i> spp.) Brown rot ( <i>Monilinia</i> spp.) Leaf spot ( <i>Blumeriella jaapii</i> ) Powdery mildew ( <i>Sphaerotheca</i> spp., <i>Podosphaera</i> spp.) Ripe fruit rot ( <i>Monilinia fructicola</i> , <i>Monilinia laxa</i> , <i>Botrytis cinerea</i> , <i>Rhizopus</i> spp.) Rust ( <i>Tranzschelia discolor</i> ) Scab ( <i>Cladosporium carpophilum</i> ) Shothole ( <i>Wilsonomyces carpophilus</i> )	10.5 to 14.5	5	72.5	0
Nectarine Peach	<b>Suppression Only:</b> Leaf curl* ( <i>Taphrina deformans</i> )				

**Application Directions.** Begin application of **Pristine** at pink bud or prior to onset of disease development and continue on a 7 to 14 day interval.

Use the shorter interval and/or the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

For aerial application to stone fruit trees, use no less than 10 gallons of spray solution per acre.

\* Not registered for use in California.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Strawberry</b>	Anthracnose <i>(Colletotrichum spp.)</i> Botrytis gray mold <i>(Botrytis cinerea)</i> Leaf spot <i>(Mycosphaerella fragariae)</i> Powdery mildew <i>(Sphaerotheca macularis)</i>	18.5 to 23	5	115	0

**Application Directions.** Begin applications of **Pristine** no later than 10% bloom, or prior to disease development, and continue on a 7 to 14 day interval.

Use the higher rate and the shorter interval when disease pressure is high.

The restricted-entry interval (REI) for treated strawberries is **12 hours**. Refer to the **Agricultural Use Requirements** box for PPE required for early entry to treated areas as permitted under the Worker Protection Standard.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Tree nut group</b> Almond Beech nut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert Hickory nut Macadamia nut Pecan Pistachio Walnut (black and English)	Alternaria leaf spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum</i> spp., <i>Marssonina juglandis</i> ) Blossom blight ( <i>Monilinia</i> spp.) Botrytis blossom and shoot blight/Green fruit rot ( <i>Botrytis cinerea</i> ) Eastern filbert blight ( <i>Anisogramma anomala</i> ) Leaf rust ( <i>Tranzschelia discolor</i> ) Panicle and shoot blight ( <i>Botryosphaeria</i> spp.) Scab ( <i>Cladosporium</i> spp.) Shothole ( <i>Wilsonomyces carpophilus</i> )	10.5 to 14.5	4	58	14 (for almond - 25 days)

**Application Directions.** In almond, begin applications of **Pristine** at pink bud and continue on a 7 to 14 day interval up to 25 days before harvest. In filbert, begin applications at budswell to budbreak, prior to infection and onset of disease development. Continue on a 7 to 14 day interval to cover and protect new growth. In pecan, begin applications of **Pristine** prior to onset of disease development and continue on a 7 to 21 day interval for the control of scab. In pistachio, begin applications prior to the onset of disease development and continue on a 10 to 30 day interval. For all other crops listed above, apply **Pristine** prior to disease development and continue on a 7 to 28 day interval. In all cases, use the shorter interval when shoot growth is very rapid.

Use the shorter interval and/or the higher rate when disease pressure is high.

No restriction on livestock feeding for almond hulls.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season.

**DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

For aerial application to tree nuts, use no less than 10 gallons of spray solution per acre.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Tropical fruit group</b> Avocado Black sapote Canistel Mamey sapote Mango Papaya Sapodilla Star apple	Anthracnose ( <i>Colletotrichum gloeosporioides</i> )  Black spot ( <i>Asperisporium caricae</i> , <i>Alternaria alternata</i> , <i>Cercospora papayae</i> )  Dry rot ( <i>Mycosphaerella</i> spp.)  Powdery mildew ( <i>Oidium</i> spp., <i>Erysiphe</i> spp.)  Pseudocercospora spot/blotch ( <i>Pseudocercospora purpurea</i> , <i>Cercospora</i> spp.)  Scab ( <i>Elsinoe mangiferae</i> )	18.5	2	37	0

**Application Directions.** Begin application of **Pristine** prior to the onset of disease development and repeat application 7 days later, as needed, or alternate with another labeled fungicide having a different mode of action.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Pristine** per season. Alternate to a labeled **non-Group 7** or **non-Group 11** fungicide with a different mode of action following two (2) sequential **Pristine** applications.

**Table 2. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Turnip greens	Alternaria leaf spot and blight <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Cercospora leaf spot <i>(Cercospora brassicicola)</i> Downy mildew <i>(Peronospora parasitica)</i> Gray mold <i>(Botrytis cinerea)</i> Powdery mildew <i>(Erysiphe polygoni)</i> Rhizoctonia stem rot and Bottom rot <i>(Rhizoctonia solani)</i> Ring spot <i>(Mycosphaerella brassicicola)</i> Sclerotinia stem rot <i>(Sclerotinia sclerotiorum, S. minor)</i> Southern blight <i>(Sclerotium rolfsii)</i> White rust <i>(Albugo candida)</i>	18.8 to 25	2	50	14

**Application Directions.** Begin **Pristine** applications prior to disease development and continue on a 7 to 10 day interval.

Use the higher rate and shorter interval when disease pressure is high.

**Resistance Management.** To limit development of resistance, **DO NOT** make more than two (2) sequential **Pristine** applications per season. Alternate to a labeled **non-Group 7** or **non-Group 11** fungicide with a different mode of action following two (2) sequential **Pristine** applications.

**Conditions of Sale and Warranty**

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Cabrio, Endura, Headline and Pristine are registered trademarks of BASF.*

*Botran is a registered trademark of Gowan Company.*

*Kelthane, Lorsban, Rally and M-Pede are registered trademarks of Dow AgroSciences LLC.*

*Lannate is a registered trademark of E.I. duPont de Nemours and Company.*

*Phaser is a registered trademark of Bayer CropScience.*

*Sonoma is a registered trademark of Albaugh, Inc.*

*Thiodan is a registered trademark of Canadian Hoechst, Ltd.*

© 2014 BASF Corporation  
All rights reserved.

007969-00199.20140613.NVA 2014-04-156-0145

Supersedes: NVA 2014-04-156-0019

Supplemental: NVA 2014-04-156-0146

Supplemental: NVA 2011-04-156-0105


BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company

45/  
47

Group	7	11	Fungicide
-------	---	----	-----------



**For the control of bud rot and *Ramularia* leaf spot in globe artichoke**

This supplemental label expires December 31, 2017 and must not be used or distributed after this date.

**Active Ingredients:**

pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester) .....	12.8%
boscalid: 3-pyridinecarboxamide,2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)- .....	25.2%
<b>Other Ingredients:</b> .....	<u>62.0%</u>
<b>Total:</b> .....	100.0%


**EPA Reg. No. 7969-199**

**Directions For Use**

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Pristine® fungicide** container label; EPA Reg. No. 7969-199, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Pristine** before applying.
- Use of **Pristine** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Pristine**.

**Application Instructions**

Refer to the **Pristine** container label for additional application instructions and restrictions.



46/47

**Pristine® fungicide Crop-specific Requirements**

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Globe artichoke	Bud rot ( <i>Botrytis cinerea</i> ) Ramularia leaf spot ( <i>Ramularia</i> spp.)	18.5 to 23.0	3	69	0

**Application Directions. Dosage and frequency/timing of applications.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7 to 14 day interval. For artichoke bud rot, begin applications at the initiation of the bud protection phase when approximately 25% of the plants have bolted. Use the shorter interval and/or the higher rate when disease pressure is high.

**DO NOT** apply **Pristine** to artichokes as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives or anything other than water. Mix **Pristine** with water only for applications to artichokes.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** exceed the specified number of applications of **Pristine** or other **Group 7** or **Group 11** fungicides per season. Adhere to the label instructions regarding the consecutive use of **Pristine** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

### Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Pristine is a registered trademark of BASF.*

©2014 BASF Corporation  
All rights reserved.

007969-00199.20140516.NVA 2014-04-156-0146

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company